

A Search for High-redshift Quasars Among GB/FIRST Flat-Spectrum Radio Sources

I. M. Hook^{1*}, R. H. Becker², R. G. McMahon³, R. L. White⁴

¹ *U.C. Berkeley Astronomy Dept, Berkeley, CA 94720, U.S.A.*

² *University of California at Davis, Davis, CA 95616, U.S.A.*

³ *Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, U.K.*

⁴ *Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, U.S.A.*

email: ihook@eso.org, bob@igpp.llnl.gov, rgm@ast.cam.ac.uk, rlw@stsci.edu

1 February 2008

ABSTRACT

We present the method and first results of a survey for high-redshift ($z > 3$) radio-loud quasars, which is based on optical identifications of 2902 flat-spectrum radio sources with $S_{5\text{GHz}} \geq 25\text{mJy}$. The radio sample was defined over a 1600 sq degree region using the 5GHz Green Bank survey and the 1.4GHz VLA FIRST survey. 560 sources were identified to a limit of $E=19.5$ on APM scans of POSS-I plates and 337 of these optical counterparts are unresolved. From these a complete sample of 73 sources for spectroscopic follow up was defined based on criteria of red ($O - E \geq 1.2$) optical colour. We have obtained spectra for 36 of these and an additional 14 had redshifts in the literature, thus 70% of the spectroscopic sample is completed. Six objects in the sample were found to be radio-loud quasars with $z > 3$ of which two were previously known. The efficiency of the spectroscopic phase of the survey is therefore about 1 in 9, whereas without the colour selection criterion the efficiency would have been 1 in ~ 40 . The six $z > 3$ quasars were found in an effective area of 1100 square degrees, implying a surface density of one flat-spectrum $z > 3$ radio-loud quasar per 190 square degrees to limits of $E=19.5$ and $S_{5\text{GHz}} \geq 25\text{mJy}$. This survey also produced the first known radio-loud BAL quasar, 1556+3517 with $z = 1.48$, which has been reported in an earlier paper (Becker et al 1997). This object has a redder optical colour ($O - E = 2.56$) than all the $z > 3$ quasars found in this survey to date. In addition we have obtained spectra of 22 GB/FIRST sources which are not part of the complete sample. We give positions, E (red) magnitudes, $O - E$ colours, radio fluxes, radio spectral index and redshifts where possible for objects for which we have obtained spectra. We give spectra and finding charts for the $z > 3$ quasars.

Key words: quasars:general radio

1 INTRODUCTION

We have begun a search for high-redshift quasars selected using the VLA FIRST survey (Becker, White & Helfand 1995). The aim of this work is to extend recent searches for radio-loud high-redshift quasars (e.g. Hook et al. 1995, 1996; Hook & McMahon 1998; Shaver et al. 1996) fainter in radio flux density by a factor of ~ 10 , and, ultimately, to use the new sample to constrain the faint end of the radio quasar luminosity function at high redshift. In addition, the sample can be used to provide an unbiased sample of damped ab-

sorption systems that can be used to study galaxy evolution at high redshift.

Whilst radio-loud quasars are only a small subset of the quasar population, the construction of radio-loud quasar samples is less prone to selection effects than are optical samples, since radio emission is unaffected by either intrinsic or extrinsic absorption due to dust. Fall & Pei (1993) discuss how dust within intervening galaxies may affect the observed evolution in optically selected samples of quasars.

Our survey technique involves optical identification of a large number of radio sources, followed by selection of the optically red, unresolved objects. The technique is similar to that used by Hook et al. (1995,1996) to find ~ 30 $z > 3$ quasars at higher radio flux densities, and complements the FIRST Bright Quasar sample (FBQS, Gregg et al. 1996).

* Present address: European Southern Observatory, Karl Schwarzschild Straße 2, D-85748 Garching b. München, Germany

The radio data are described in section 2 of this paper and the optical identification of the radio sample and selection of the spectroscopic sample are described in section 3. The spectroscopic observations are described in section 4, and the results are presented in section 5. Finally, tables of redshifts, optical and radio data are given for the objects for which we have obtained spectra, along with the spectra and finding charts for the $z > 3$ quasars.

2 DEFINITION OF THE RADIO SAMPLE

This study uses as its starting point the 5GHz 87GB catalog of Gregory & Condon (1991). By defining the sample at high frequency, flat-spectrum core-dominated objects, which are usually identified with quasars, are preferentially selected. Spectral index information and the accurate positions (better than $\pm 1.0''$) needed for optical identification are provided by the $S_{1.4\text{GHz}} \geq 1\text{mJy}$ VLA ‘FIRST’ survey (Becker, White & Helfand, 1995). The sample covers an area of 0.49sr (1600sq deg), limited by the extent of the FIRST survey which at the commencement of this work covered the region $7^{\text{h}}30^{\text{m}} < \alpha(\text{J2000}) < 17^{\text{h}}30^{\text{m}}$, $28^{\circ}30' < \delta(\text{J2000}) < 41^{\circ}30'$.

Objects with $S > 25\text{mJy}$ from the 5GHz 87GB catalog were matched to the FIRST catalog with a matching radius of 1 arcmin. In cases where more than one FIRST object matched the 5GHz source (about 50% of the time), the FIRST source with the strongest flux was selected (i.e. the source with the steepest spectrum). The reason for this is that if that source then passes the flat spectrum selection criterion then there is certain to be a flat spectrum source associated with the 5GHz source.

Matched sources with radio spectral index flatter than $\alpha_{1.4\text{GHz}}^{5\text{GHz}} = -0.5$ (where $S \propto \nu^\alpha$) were then selected. The exact spectral index criterion chosen was somewhat arbitrary but this has proved successful in the past at separating the quasar and radio galaxy populations. The ‘total’ (rather than ‘peak’) fluxes from the 1.4GHz FIRST catalogue were used to compute the spectral index. A total of 2902 sources ($\sim 2\text{deg}^{-2}$) meet these criteria.

3 OPTICAL IDENTIFICATION AND SELECTION OF THE SPECTROSCOPIC SAMPLE

Optical identification of the radio sources was carried out using APM (Automated Plate Measurement Facility at Cambridge, U.K.) scans of POSS-I E (red) and O (blue) plates. The identification procedure and selection of candidates is similar to that in Hook et al. (1996), and more details can be found there along with a description of the APM POSS-I catalog.

Identifications were made based on positional coincidence of the FIRST radio position with an optical counterpart on the POSS-I E plate. The sample of candidate high-redshift quasars was then selected using the APM image classification, $N\sigma_c$, measured from the E plate, and the O–E colour. The basis for the colour selection method is that the O – E colours of quasars with $z > 2$ become rapidly redder with redshift, due to absorption by intervening Ly α (see figure 1 in Hook et al. 1995). To define a sample with

a high level of completeness at $z > 3$ we chose a limit of $O - E \geq 1.2$. A magnitude limit of $E \leq 19.5$ was imposed (0.5 mag brighter than the E plate limit) so that spectroscopy with the Lick-3m would be more feasible. The APM star/galaxy separation is also more reliable away from the plate limit.

Based on results from the $S \geq 200\text{mJy}$ high-redshift radio-quasar survey of Hook et al. (1996) and on results from the FBQS (Gregg et al. 1996), stricter selection criteria were used for this new survey. The criterion for positional coincidence used by Hook et al. (1996) was $\Delta r \leq 3.0''$ and for the FBQS it was $\Delta r \leq 2.0''$ (Gregg et al. 1996). Even that was found to be too generous: 64 out of 69 (93%) spectroscopically confirmed quasars in FBQS were found to have $\Delta r \leq 1.0''$, and 97% have $\Delta r \leq 1.1''$ (the outliers were extended radio sources). Since the current high redshift survey goes 2 magnitudes fainter in E than FBQS, we adopt a more conservative criterion of $\Delta r \leq 1.5''$. Figure 1 shows this limit in relation to the histogram of Δr for identifications with $E \leq 19.5$. The distribution peaks for the stellar and non-stellar identifications at $\Delta r \sim 0.5''$. However, whereas beyond $\sim 3''$ the stellar histogram is indistinguishable from the derived background, the non-stellar identifications show an excess. There is evidence from Figure 1a that this may persist out to a radius of $\sim 40''$ (and possibly further, although this would not be apparent in Fig 1a since the background was determined by fitting the distribution from $40''$ to $100''$). This may be due to either the clustering of galaxies or that the peak of the radio emission is not on the optical counterpart as would be expected for extended asymmetric radio sources. Since the extended or resolved optical counterparts are not the subject of this paper, we do not consider this issue any further. This issue is discussed further in a forthcoming paper (McMahon et al, in prep) on the optical identifications of all FIRST sources using the APM POSS-I catalogue.

There is a small excess of 34 stellar objects above the background level between radii of $1.5''$ and $3.0''$ (see Fig 1b). If we assume that these are real identifications that are excluded from the spectroscopic sample, then the incompleteness due to the $\Delta r \leq 1.5''$ criterion is $\sim 9\%$. This agrees with the fact that of the 13 $z > 3$ quasars found in the survey of Hook et al. (1996) (with limiting magnitude $E \leq 20$), only one would not have satisfied the criterion of $\Delta r \leq 1.5''$.

The background counts for stellar sources with $E \leq 19.5$ was found to be 0.64 per square arcmin (see Fig. 1 and caption). Thus within a search radius of $1.5''$ we expect 3.6 false identifications amongst the stellar identifications, i.e. 1.1%. The corresponding background for red ($O - E \geq 1.2$) stellar sources is 0.54 per square arcmin, and we therefore expect 3.1 false red stellar identifications. In section 5 this number is compared to the number of stars found in our sample.

We also now adopt a stricter criterion for stellar image classification. In Hook et al. (1996) a criterion of $|N\sigma_c| \leq 3.0$ was used (where $N\sigma_c$ can be considered as the number of standard deviations a particular object is from having a stellar profile; see Hook et al. (1996) for a more detailed explanation). However only one of the 13 $z > 3$ quasars found in that survey had $|N\sigma_c| > 2.0$, an object with $E=19.93$, fainter than the limit of the current survey. Thus we now adopt a criterion of $|N\sigma_c| \leq 2.0$. Whilst using these stricter




Figure 1. Histograms of positional difference Δr between the optical position measured by the APM and the FIRST radio position for the flat spectrum sample. (b) shows a blowup of the region $\Delta r \leq 10.0''$. The open histogram shows all identifications with $E \leq 19.5$ and the shaded histogram shows those which are classified as stellar ($N\sigma_c \leq 2.0$, see text). The dashed line shows the limit used to define the spectroscopic sample, $\Delta r \leq 1.5''$. The solid sloping lines show fits to the histograms in the range $40.0'' < \Delta r < 100.0''$, which determine the background counts. The upper line corresponds to 1.03 background sources with $E \leq 19.5$ per arcmin². The lower line corresponds to 0.64 stellar sources per arcmin² to the same magnitude limit.

criteria on Δr and $N\sigma_c$ has little effect on the completeness of the final high-redshift quasar sample compared to that in Hook et al. (1996), the number of candidates requiring spectroscopy is reduced by $\sim 40\%$.

In summary the selection criteria are (i) $S_{5\text{GHz}} \geq 25\text{mJy}$, (ii) flat radio spectral index $\alpha_{1.4\text{GHz}}^{5\text{GHz}} \geq -0.5$, (iii) positional coincidence, $\Delta r \leq 1.5''$, (iv) red optical colour $O - E \geq 1.2$, (v) unresolved image on the E plate, $|N\sigma_c| \leq 2.0$, (vi) $E \leq 19.5$ mag. The optical counterpart was not required to be detected on the O plate.

There is little overlap between this sample and the FBQS. The FBQS does not have a radio spectral index cut and reaches $S_{1.4\text{GHz}} \sim 1\text{mJy}$. FBQS now has a blue optical colour criterion of $O - E < 2.0$ and has a significantly brighter optical limit of $E=17.5$. Hence the two quasar samples are largely complementary.

Of the 2902 flat spectrum sources (as defined in section 2) in the GB/FIRST survey area, 560 have optical identifications brighter than $E=19.5$ within $1.5''$. The histogram of their $O - E$ colours, shown in Figure 2, has a bimodal distribution. The bluer peak is dominated by stellar identifications (the vast majority of which are low-redshift quasars) and the redder peak is dominated by galaxies. By imposing a red colour selection criterion on the sample of stellar identifications, the majority of low redshift quasars are eliminated. Of the 560 identifications, 337 are classified as stellar and 73 of these satisfy the colour criterion above ($O - E \geq 1.2$). This sample of 73 objects will be referred to as the ‘spectroscopic sample’ and they are shown in the colour-magnitude diagram in Figure 3.

Fourteen of the spectroscopic sample had published spectroscopic data, leaving 59 objects requiring spectroscopy. The previously-known objects were not re-




Figure 2. Colour histogram for APM identifications of flat spectrum GB/FIRST sources. The open histogram shows all identifications (both stars and galaxies) within $1.5''$ and with $E \leq 19.5$. The hatched histogram shows the stellar sources only (337) and the solid histogram shows the stellar sources which were not detected on the O plate and for which the $O - E$ values are limits. The dashed line shows the colour selection criterion adopted ($O - E \geq 1.2$), which eliminates the blue stellar identifications (i.e. low-redshift quasars) from the spectroscopic sample.




Figure 3. Colour-magnitude diagram for APM identifications of flat-spectrum GB/FIRST sources. The circles represent objects in the spectroscopic sample: those with spectroscopic data are shown as filled circles while those with no spectra as yet are shown as open circles. Arrows show limits on $O - E$ colour for objects that were not detected on the O plate. The $z > 3$ quasars are shown by stars and the radio-loud BAL quasar 1556+3517 is shown by the square. The dashed lines show the E and $O - E$ limits of the spectroscopic sample. The extra objects observed which do not satisfy all the selection criteria (see section 5) are shown as crosses.

observed and their properties are summarised in Table 1. Two of the previously-known objects are radio-selected $z > 3$ quasars (0933+2845, Gregg et al. 1996 and 1340+3754, Hook et al. 1995).

4 OPTICAL SPECTROSCOPY

The aim of the spectroscopic phase of this project is to obtain classification (high-redshift quasar or not) for the complete spectroscopic sample defined above. This does not require redshifts to be obtained for all the objects, for example many ‘featureless’ spectra could be rejected as possible $z > 3$ quasars by the lack of $\text{Ly}\alpha$ forest absorption. The strategy was therefore to take short exposures at low dispersion to obtain spectra of sufficient quality for the nature of the object and, where possible, the redshift to be determined.

Most of the spectroscopy was carried out in 1995 April and 1996 May using the KAST double-spectrograph at the Cassegrain focus of the 3m Shane Telescope at Lick observatory. Typical exposure times were 600-900s. For the May 1996 run, a grating giving $4.6\text{\AA}/\text{pix}$ dispersion was used on the red side, and on the blue side the selected grism gave a dispersion of $2.54\text{\AA}/\text{pix}$. A dichroic at $\sim 5500\text{\AA}$ was used. The set-up was such that the blue side of the spectrum covered $2900 - 6000\text{\AA}$ and the red side covered $5080 - 10610\text{\AA}$. For the Apr 1995 run a grating giving $2.32\text{\AA}/\text{pix}$ dispersion was used on the red side, and on the blue side the selected grism gave a dispersion of $1.85\text{\AA}/\text{pix}$. A dichroic at $\sim 5500\text{\AA}$




Figure 4. Histogram of redshifts obtained. The white histogram represents the spectroscopic sample, including previously-known objects and objects with uncertain redshifts, a total of 33 objects (the two stars are not included here). Objects in the spectroscopic sample with spectra from which redshifts could not be measured (15 objects - it is unlikely that any of these have $z > 3$; see text for explanation) are represented by the white box and objects for which spectra have not been obtained (23) are represented by the hatched box. Similarly the black histogram and rectangle show the extra objects observed which satisfy a less stringent set of selection criteria described in section 5 (15 of these have definite redshifts and 7 have ‘featureless’ spectra).

was used giving a total wavelength coverage of $3430-8150\text{\AA}$. For both runs the detectors on both the red and blue sides were 1200×400 UV-flooded Reticon CCDs with 27 micron pixels, corresponding to about $0.8 \text{ arcsec}/\text{pixel}$. The CCD chips were windowed in the spatial direction to reduce the readout time.

Seven objects from the spectroscopic sample were observed at the Keck telescope with the Low Resolution Imaging Spectrograph (LRIS). These observations were carried out as a backup program during a run to obtain spectra of distant supernovae. A 300 l/mm grating was used on LRIS and covered the wavelength range $4900 - 9900\text{\AA}$ with resolution of $2.5\text{\AA}/\text{pix}$. In addition, one object was observed at 4.2m WHT, La Palma in 1993 April and another by R. Weymann at Palomar 5m in 1991 December as part of other programs.

For all observations the width of the slit was adjusted to be compatible with the seeing at the time of observation ($1 - 2''$) and the data were taken with a long slit at the parallactic angle. Some of the objects were too faint to be easily visible on the acquisition TV at the Lick 3m so accurate offsetting from nearby stars was used to position the target object in the slit. Spectrophotometric standards were observed to calibrate the spectra.

Table 1. Optical and radio data for objects in the spectroscopic sample which had spectroscopic data in the literature.

α	Optical Position		δ	z	$N\sigma_c$	E mag	O–E mag	Δr "	S_{GB} mJy	S_{B91} mJy	S_{FIRST} mJy	S_{NVSS} mJy	$\alpha_{1.4GHz}^{5GHz}$	Reference/ Classification
09 11	47.71	+33	49 17.6	—	1.75	18.62	1.56	0.78	250	207	391.2	380.8	−0.35	H96 / featureless
09 30	55.28	+35	03 36.9	—	0.12	19.18	1.24	0.85	383	381	518.8	484.5	−0.24	H96 / featureless
09 33	37.31	+28	45 32.3	3.42	0.27	17.50	1.27	0.16	66	63	120.9	117.0	−0.48	G96 FBQS / QSO
10 50	58.15	+34	30 10.8	2.52	1.43	18.98	1.40	0.30	298	291	547.3	554.0	−0.48	W84 / QSO
11 49	00.29	+29	58 41.0	0.15	1.15	17.05	2.72	0.34	41	39	53.2	52.5	−0.20	G96 FBQS / Gal
12 35	05.79	+36	21 18.7	1.60	−0.20	19.06	1.26	0.47	250	251	175.7	154.5	0.28	H96 / QSO
13 24	12.09	+40	48 12.4	0.50	0.16	19.49	2.56	0.87	413	423	369.4	350.1	0.09	V96 / QSO
13 40	22.94	+37	54 44.4	3.10	0.08	17.85	1.68	0.85	305	301	284.3	273.9	0.06	H95 / QSO
13 50	35.95	+33	42 17.2	0.01	−0.85	11.41	1.35	0.93	76	90	100.5	103.8 ^a	−0.22	HGC95 / Gal
14 04	14.63	+28	46 36.5	—	−0.84	16.03	1.49	1.03	38	36	12.5	11.7	0.87	G96 FBQS / star
14 21	25.62	+39	43 29.4	0.62	1.43	18.33	> 3.65	0.79	67	64	43.6	82.9 ^b	0.34	V90 / QSO
14 58	44.88	+37	20 21.6	0.33	−0.67	18.22	1.70	0.68	591	591	271.9	215.2	0.61	V96, H96 / Gal
15 58	55.21	+33	23 19.3	1.65	−0.13	16.49	1.24	0.83	88	86	145.5	142.7	−0.39	F86 / QSO
16 33	02.12	+39	24 27.4	1.02	0.25	15.02	1.21	0.34	40	38	54.7	79.1	−0.25	CCH90 / QSO

Notes on column headings: S_{GB} is the 5GHz flux from Gregory & Condon (1991) and S_{B91} is the 5GHz flux from the Becker et al (1991) catalog. S_{NVSS} was determined by summing up the fluxes from sources in the 1.4GHz NRAO-VLA Sky Survey catalog (NVSS, Condon et al 1994) within 60'' of the FIRST position. S_{FIRST} is the 'Total' flux from the 1.4GHz FIRST catalog and $\alpha_{1.4GHz}^{5GHz}$ is the spectral index between S_{FIRST} and S_{GB} .

^a NVSS flux is the sum of two components. The central component has $S=97.3$ mJy and a second component 20'' away has $S=6.5$ mJy.

^b NVSS flux is the sum of two components: The closest source has $S=62.3$ mJy and a second, probably unrelated source 42'' away has $S=20.6$ mJy.

References: F86=Foltz et al (1986), G96 = Gregg et al (1996), H95 = Hook et al (1995), H96 = Hook et al (1996), HGC95= Huchra, Geller & Corwin (1995), V90=Vigotti et al (1990), V96 = Vermeulen et al (1996), W84=Wampler et al 1984, CCH90=Crampton, Cowley & Hartwick (1990)

The reduction of data was carried out using standard software from the IRAF[†] package.

5 RESULTS AND DISCUSSION

Of the 59 objects requiring spectroscopy we now present new spectroscopic data for 36. The properties of objects in the spectroscopic sample for which we obtained spectra are given in Table 2. When combined with the 14 objects which have spectroscopic data in the literature, 50 (70%) of the complete sample defined in Section 3 now have spectroscopic data. The remaining 23 objects have yet to be observed.

Four new quasars with $z > 3$ have been found in the spectroscopic sample to date. Since 36 objects were observed, the efficiency of the survey for finding $z > 3$ quasars is approximately 1 in 9, similar to that of Hook et al 1996 (although note that the selection criteria are now stricter). If we had not used the colour selection criterion, the efficiency would be lower by a factor of ~ 4.5 . The $z > 3$ quasars are marked on the colour-magnitude diagram (Figure 3) with stars. A further 8 new quasars in the range $2 < z < 3$ were found (but note that the colour selection criterion means the sample is substantially incomplete for $z < 3$).

[†] IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

The remaining observed objects were found to be low-redshift ($z < 2$) quasars, emission- or absorption-line galaxies and objects with featureless spectra which could be ruled out as possible high-redshift quasars. Two objects were found to be stars and are probably chance coincidences with the radio position (both have $\Delta r > 1.0''$). This agrees well with the number of false identifications expected (see section 3). We expect 3.1 false, red, stellar identifications in the full spectroscopic sample, thus after observing 70% of these objects we expect to have found 2.1 stars.

The first radio-loud BAL quasar, 1556+3517 with $z = 1.48$, was discovered in this survey and was reported in an earlier paper (Becker et al. 1997). Note that this object is redder ($O - E = 2.56$) than any of the $z > 3$ quasars in the sample so far.

Of the 50 objects with spectroscopic data, 35 have a definite redshift based on emission or absorption features (including the two stars). Their redshift distribution is shown in Figure 4. The 15 'featureless' objects are represented by the open box, while objects with no spectral information are represented by the hatched box.

In addition we have observed 22 objects that do not meet one or more of the selection criteria of the spectroscopic sample, but which satisfy a less stringent set of selection criteria: $\Delta r \leq 3.0''$; $O - E \geq 1.0$; $|N\sigma_c| \leq 3.0$; $E \leq 20$ mag. None of them was found to have $z > 3$. The results for these objects are summarised in Table 3. They are plotted as crosses on the colour-magnitude diagram in Figure 3 and represented by the solid shading in the redshift histogram in Figure 4.

Table 2. Optical and radio data for objects from the complete sample observed spectroscopically. These objects satisfy the selection criteria (i) $S_{5\text{GHz,GB}} \geq 25\text{mJy}$, (ii) flat radio spectral index $\alpha_{1.4\text{GHz}}^{5\text{GHz}} \geq -0.5$, (iii) positional coincidence, $\Delta r \leq 1.5''$, (iv) red optical colour $O - E \geq 1.2$, (v) unresolved image on the E plate, $|N\sigma_c| \leq 2.0$, (vi) $E \leq 19.5$ mag.

Optical Position			z	$N\sigma_c$	E	O–E	Δr	S_{GB}	S_{B91}	S_{FIRST}	S_{NVSS}	$\alpha_{1.4\text{GHz}}^{5\text{GHz}}$	Tel/Date/Class
α	J2000		δ		mag	mag	"	mJy	mJy	mJy	mJy		
08 47 15.16	+38 31 10.2	3.18	−1.01	17.55	1.37	0.31	124	122	185.8	201.1	−0.32	K 03/97 QSO	
08 52 44.76	+34 35 40.8	1.65	−1.31	18.43	1.56	0.53	70	66	69.6	69.1 ^a	0.01	K 03/97 QSO	
08 57 41.88	+38 16 35.6	0.43	1.15	19.40	> 2.63	0.47	49	47	33.8	70.9 ^b	0.29	K 03/97 Gal (abs.)	
08 59 17.31	+34 09 09.3	0.55	1.81	19.46	> 2.57	0.65	30		55.6	53.8	−0.48	K 03/97 Gal (abs.)	
09 02 39.91	+39 57 26.7	3.08	0.83	17.56	1.24	0.14	31	29	32.4		−0.03	K 03/97 QSO	
10 23 24.04	+28 56 50.8	—	1.63	19.12	1.80	0.24	89	86	127.4	139.7	−0.28	W 04/93 featureless	
10 28 46.95	+41 26 56.6	2.82*	1.11	19.46	1.50	0.19	53	51	62.3	74.4	−0.13	L 05/96 QSO ?	
10 42 06.79	+29 38 26.8	2.55	−0.09	19.30	1.21	0.73	56	54	70.0	71.9	−0.18	L 04/95 QSO	
10 42 32.27	+36 15 20.7	—	1.81	18.62	1.60	0.09	92	89	60.1	68.1	0.33	L 05/96 featureless ?	
10 44 06.39	+29 59 01.3	2.99	0.61	18.62	1.26	0.73	80	79	37.4	40.9	0.60	P 12/91 QSO	
10 55 38.67	+30 52 50.6	—	1.69	17.83	2.27	0.65	32	31	7.8	14.7	1.11	L 04/95 not QSO	
11 21 11.12	+37 02 47.0	1.62	−1.09	17.04	1.22	0.16	72	69	100.8	105.7	−0.26	L 05/96 QSO	
11 29 14.28	+37 03 17.5	—	−0.99	18.34	1.43	0.52	95	91	70.1	60.9	0.24	L 05/96 featureless	
11 51 16.95	+40 08 21.6	2.74	−0.88	19.41	> 2.87	0.52	111	108	84.0	81.7	0.22	L 05/96 QSO	
12 34 44.34	+36 37 22.2	—	−1.07	19.10	> 2.67	0.27	45	43	62.8	64.3	−0.26	L 05/96 Gal (abs?)	
12 36 23.02	+39 00 00.7	—	0.01	17.32	1.36	0.32	47	57	37.0	37.1	0.19	L 05/96 featureless	
12 58 32.18	+29 09 03.0	3.47	0.32	18.52	2.06	0.48	32	32	25.0	25.1	0.19	L 04/95 QSO	
13 01 23.40	+28 42 53.8	0.42	1.40	18.86	> 3.26	0.47	31		13.1	12.1	0.68	L 04/95 Gal (em.)	
13 24 04.90	+34 17 38.5	—	1.64	18.80	2.45	0.35	85	83	54.8	62.2	0.34	L 05/96 featureless ?	
13 42 52.97	+40 32 02.0	0.91	0.99	19.50	1.45	0.58	113	99	137.2	152.2	−0.15	L 05/96 QSO	
13 48 48.13	+29 39 17.6	0.24	2.00	18.43	2.80	1.19	216	213	327.5	695.6	−0.33	L 04/95 Gal	
13 58 19.85	+32 19 27.8	2.85	−0.27	18.66	> 3.09	1.32	36	36	20.6	18.8	0.44	L 05/96 QSO	
14 01 38.82	+36 11 21.6	—	0.60	19.03	1.95	1.29	61	59	83.3	73.0	−0.24	L 05/96 featureless	
14 13 14.15	+32 00 56.6	—	1.80	19.05	2.89	1.19	31		44.0	49.0	−0.28	L 05/96 featureless	
14 36 42.84	+40 22 05.8	2.57	−0.93	18.07	1.21	0.19	33	32	23.0	22.3	0.28	L 05/96 QSO	
14 53 18.56	+35 05 39.4	—	−0.13	19.48	1.71	0.51	127	123	215.8	217.3	−0.42	L 05/96 featureless	
15 23 14.87	+38 14 01.7	3.15	−1.09	18.38	1.40	0.40	27		47.7	52.6	−0.45	L 05/96 QSO	
15 29 42.17	+35 08 50.7	0.29	0.85	17.83	2.18	0.94	50	48	90.0	105.6	−0.46	L 05/96 QSO	
15 42 46.77	+33 46 02.4	0.28	1.99	17.85	3.27	0.93	69	66	36.8	39.8	0.49	L 05/96 Gal	
15 56 33.79	+35 17 56.9	1.48	−1.71	18.68	2.56	0.71	27		31.8	28.1	−0.13	L 05/96 BAL QSO	
16 05 23.73	+30 38 37.0	2.67	0.27	18.66	> 3.51	0.44	76	59	52.6	50.0	0.29	L 04/95 QSO	
16 20 04.77	+35 15 54.5	2.95	0.91	17.94	1.32	1.01	40	38	50.1	57.9	−0.18	L 05/96 QSO	
16 55 42.82	+32 44 20.0	—	−1.37	17.84	1.35	0.30	69	69	31.8	38.7	0.61	L 05/96 featureless ?	
17 00 08.72	+29 19 03.1	—	−2.00	16.02	1.41	1.08	39	48	3.0	5.1	2.01	L 05/96 star	
17 01 24.70	+39 54 37.3	—	0.64	16.58	1.31	0.82	159	157	256.8	190.8	−0.38	L 05/96 featureless	
17 06 50.43	+30 04 12.1	—	−0.03	18.25	1.31	0.32	38	36	27.9	29.5	0.24	L 05/96 featureless	

Notes

* uncertain redshift.

Telescope code: L=Lick 3m, K=Keck-II, W=WHT, P=Palomar 5m (Weymann, private communication).

^a NVSS flux contains contributions from two components: a central component with flux density $S=60.7\text{mJy}$ and another $14''$ away with $S=8.4\text{mJy}$.

^b As above but central component has $S=54.5\text{mJy}$ and other component $36''$ away has $S=16.2\text{mJy}$.

5.1 The new sample of $z > 3$ quasars

A summary of the six $z > 3$ quasars in the sample to date is given in Table 4, their spectra are shown in Figure 5 and finding charts are given in Figure 6. Note that the spectra obtained at Keck (0847+3831 and 0902+3957) have been corrected for atmospheric absorption bands while the others have not. The spectrum of 0933+2845 ($z = 3.42$, Gregg et al 1996) was obtained at Lick on 1995 July 3 as part of the FBQS. 1258+2909 was re-observed at Lick 3m on 1996 March 15 to obtain a higher signal-to-noise spectrum. This spectrum, rather than the discovery spectrum, is shown in Figure 5.

Two $z > 3$ radio-loud quasars are known to lie within

the survey area but were not found by this survey. FBQS J1021+3001 with $z = 3.1$ (Gregg et al 1996) is in the FIRST catalogue but is below the flux limit used in this search. B3 1239+376 with $z = 3.82$ (Vermeulen et al. 1996) is optically fainter than the limit of the APM scans.

6 CONCLUSIONS

In this paper we have described a survey for high redshift radio-loud quasars with a limiting flux density of 25mJy , about a factor of 10 fainter than previous surveys. This survey has become feasible because of new accurate positions for large numbers of faint radio sources supplied by the VLA




Figure 5. Spectra of the $z > 3$ quasars in the GB/FIRST flat-spectrum sample to date.

Table 3. Optical and radio data for objects observed spectroscopically that were not part of the complete spectroscopic sample. These objects satisfy a less stringent set of selection criteria: $\Delta r \leq 3.0''$; $O - E \geq 1.0$; $|N\sigma_c| \leq 3.0$; $E \leq 20$ mag.

α	Optical Position		δ	z	$N\sigma_c$	E	O-E	Δr	S_{GB}	S_{B91}	S_{FIRST}	S_{NVSS}	$\alpha_{1.4GHz}^{5GHz}$	Tel/Date/Class
	J2000					mag	mag	"	mJy	mJy	mJy	mJy		
08 35	19.76	+29 09	21.2	1.27	1.37	19.72	2.14	0.55	36	45	41.3	38.9	-0.11	L 04/95 QSO
09 00	30.44	+33 30	09.7	—	1.09	19.91	1.25	0.86	81	77	72.4	82.5	0.09	K 03/97 featureless
09 06	02.49	+41 16	27.6	0.41	3.00	17.85	> 3.82	2.75	279	276	771.1	848.3	-0.80	K 03/97 QSO
09 08	53.08	+30 47	38.3	1.91	0.65	19.63	1.08	0.27	35	32	34.2	40.1	0.02	L 04/95 QSO
09 24	38.22	+30 28	36.6	0.27	2.95	17.76	3.11	0.70	27		4.0	8.4	1.50	L 04/95 Gal
09 27	39.78	+30 34	17.3	1.20	-0.24	19.59	1.41	0.87	57	56	46.1	51.5	0.17	L 04/95 QSO
11 05	50.05	+30 01	46.2	—	2.07	19.36	1.39	0.79	53	50	64.2		-0.15	L 04/95 featureless
11 24	29.65	+28 31	25.3	1.35	1.08	19.04	1.10	0.46	31	38	22.3	24.0	0.26	L 04/95 QSO
11 27	29.21	+37 56	12.4	—	-0.09	19.60	> 2.00	1.13	70	67	66.4	78.3	0.04	L 05/96 featureless ?
12 11	34.02	+28 47	33.1	—	-0.81	19.65	> 2.26	0.17	35	33	32.9	32.0	0.05	L 04/95 ?
12 24	09.80	+28 35	10.3	0.40	0.89	19.82	2.06	0.84	77	74	18.6	297.8 ^a	1.12	L 04/95 Gal (em.)
12 39	14.99	+40 49	55.2	1.31	-0.37	17.55	1.18	1.02	78	74	65.7	70.3	0.13	L 05/96 QSO
13 01	56.57	+29 04	55.7	2.29	-0.57	19.19	1.10	0.44	50	48	97.5	103.1	-0.52	L 04/95 QSO
13 24	57.44	+32 51	46.2	1.74	0.11	17.91	1.01	1.21	92	92	57.6	56.7	0.37	L 05/96 QSO
13 26	13.76	+28 51	43.6	1.29	1.25	18.14	1.04	0.69	31	46	20.9	36.8	0.31	L 04/95 QSO
14 08	35.27	+36 17	59.7	—	-1.09	18.28	1.01	0.31	82	79	128.3	133.3	-0.35	L 05/96 featureless
14 13	13.65	+41 16	37.8	2.61	1.41	17.71	1.06	0.34	35	33	26.7	30.3	0.21	L 05/96 QSO
14 36	27.24	+41 29	31.6	0.17*	1.01	18.77	2.09	1.63	41	38	45.6	53.6	-0.08	L 05/96 Gal
14 45	27.10	+30 51	28.6	0.42	2.51	19.01	2.84	0.57	35	107	6.3	12.7	1.35	L 04/95 Gal (em.)
15 36	27.97	+37 42	05.9	—	-0.41	19.53	> 1.99	1.00	41	49	47.0		-0.11	L 05/95 featureless
17 03	22.63	+38 59	29.0	1.08	1.73	18.33	1.25	1.56	52	49	97.4	104.9	-0.49	L 05/96 QSO
17 07	35.97	+35 39	50.0	—	2.13	16.66	2.71	0.51	53	51	100.1		-0.50	L 05/96 featureless

* uncertain redshift.

^a The source has complex structure in the FIRST map. The FIRST flux given is the flux of the central component only. The NVSS flux is the sum of 4 components within $60''$, the most central of which has $S=110.6$ mJy.**Table 4.** Summary of the $z > 3$ QSOs in the sample to date.

Name ¹	z	$N\sigma_c$	E	O - E	$\Delta r/''$	S_{GB}/mJy	$\alpha_{1.4GHz}^{5GHz}$	Comments
0847+3831	3.18	-1.01	17.55	1.37	0.31	124	-0.32	Keck 03/97
0902+3957	3.08	0.83	17.56	1.24	0.14	31	-0.03	Keck 03/97
0933+2845	3.42	0.27	17.50	1.27	0.16	66	-0.48	Previously known, Gregg et al 1996
1258+2909	3.47	0.32	18.52	2.06	0.48	32	0.19	Lick 04/95
1340+3754	3.10	0.08	17.85	1.68	0.85	305	0.06	Previously known, Hook et al 1995
1523+3814	3.15	-1.09	18.38	1.40	0.40	27	-0.45	Lick 05/96

Notes¹ based on J2000 position

FIRST survey. Six $z > 3$ quasars have so far been found in an area of $0.49sr$ (~ 1610 sq deg), after completing follow up for 70% of the spectroscopic sample. The effective area covered so far is thus $70\% \times 1610 = 1130$ square degrees. Therefore we find a surface density of flat-spectrum $z > 3$ quasars of one per 190 square degrees to a radio flux limit of $S_{5GHz} \geq 25$ mJy and an optical limit of $E=19.5$ mag. After completion of the spectroscopic observations the new sample will be used in an analysis of the space density of quasars at high redshift.

ACKNOWLEDGMENTS

IMH acknowledges a PPARC/NATO fellowship and RGM thanks the Royal Society for support. This research has

made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration. Some observations described in this paper were obtained at the Keck-II 10m telescope of the California Association for Research in Astronomy. We thank Daniel Stern, Lisa Storrie-Lombardi, Robert Knop and Saul Perlmutter for assistance with the observations.

REFERENCES

- Becker R. H., Gregg M. D., Hook I. M., McMahon R. G., White R. L., Helfand D. J., 1997, ApJ, 479, L93
 Becker R. H., White R. L., Edwards A. L., 1991, ApJS, 75, 1
 Becker R. H., White R. L., Helfand D. J., 1995, ApJ, 450, 559
 Condon J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A.,




Figure 6. Finding charts from APM scans of POSS-I E plates ($\sim R$ band) for the $z > 3$ quasars in the GB/FIRST flat-spectrum sample to date. North is at the top, East to the left. The central cross is 1 arcmin across.

- Broderick J. J., 1994, in *Astronomical Data Analysis Software & Systems III*, ASP, ed. Crabtree D. R. et al., 155
- Crampton D., Cowley A. P., Hartwick F. D. A., 1990, *AJ*, 100, 47
- Fall S.M., Pei Y., 1993, *ApJ*, 402, 479
- Foltz C., Weymann R., Peterson B. M., Sun L., Malkan M. A., Chaffee F. H., 1986, *ApJ*, 307, 504
- Gregory P.C., Condon J.J., 1991, *ApJS*, 75, 1011
- Gregg M. D., Becker R. H., White R. L., Helfand D. J., McMahon R. G., Hook I. M., 1996, *AJ*, 112, 407
- Hook I.M., McMahon R.G., Irwin, M. J., Hazard, C., 1996, *MNRAS*, 282, 1274
- Hook I.M., McMahon R.G., Patnaik A.R., Browne I.W.A., Wilkinson P.N., Irwin M.J., Hazard C., 1995, *MNRAS*, 273, L63
- Hook I.M., McMahon R.G., 1998, *MNRAS*, 294, L7
- Huchra J.P., Geller M. J., Corwin H. G., 1995, *ApJS*, 99, 391
- Shaver P.A., Wall J.V., Kellermann K.I., Jackson C.A., Hawkins M.R.S., 1996, *Nat* 384, 439
- Vermeulen R., Taylor, G. B., Readhead, A. C. S., Browne, I. W. A., 1996, *AJ*, 111, 1013
- Vigotti M., Vettolani G., Merighi R., Lahulla J. F., Lopez-Arroyo M., 1990, *A&ASS*, 83, 205
- Wampler E. J., Gaskell, C. M., Burke W. L., Baldwin J. A., 1984, *ApJ*, 276, 403